

Success factors in male WTF taekwondo juniors

Jerzy Sadowski¹, Dariusz Gierczuk¹, Jerzy Miller², Igor Cieśliński¹, Mariusz Buszta¹

¹ Faculty of Physical Education and Sport in Biała Podlaska, Poland

² The State College of Computer Science and Business Administration in Łomża, Poland

Key words: WTF taekwondo, regression model, complex reaction, technical factors

Summary

Introduction. The aim of the study was to identify selected success factors of elite Olympic taekwondo competitors in the context of medals that they have or have not won during Polish Junior Championships (PJCcs).

Material and methods. Sixty-three elite male taekwondo competitors took part in the study. The competitors were divided into two groups on the basis of their achievements: PJC medallists and non-medallists. Factors determining sports success were modelled with the use of a general linear model (GLM). The procedure lasted until the minimum of the quality criterion, i.e., the Akaike information criterion (AIC), was reached.

Results. Successful competitors (medallists) demonstrated significantly better conditioning, technique, coordination and structural parameters. Dynamic strength, general endurance, technical skills (e.g., roundhouse kicks), complex reactions and low adiposity levels were the factors that contributed to the achievement of excellent results in taekwondo.

Conclusions. The research results revealed groups of success factors in taekwondo at a national level. Variables concerning coordination factors (complex reaction – 16%) and technical factors (360° dollyo chagi montong – 8.8%) exerted the greatest influence on point scores.

Introduction

The most important success factors in sports include body build, conditioning, technical and tactical skills, mental abilities and the experience of a competitor [1]. Since WTF (World Taekwondo Federation) taekwondo was approved for the Sydney 2000 Olympic Games, it has been the subject of extensive research, where one of the areas of interest is the identification of success factors. Training effectiveness in sport is measured with the best possible performance (result) in the most important competitions. Therefore, many researchers have been interested in establishing success factors in taekwondo.

The majority of studies have sought to determine correlations between different indicators and performance in taekwondo, including morphological [2-10], biomechanical [7; 11], conditioning, physiological and psychological indicators [6; 8; 12-14] and indicators related to coordination [15]. The aforementioned researchers have found that taekwondo competitors with mesomorphic body build, high levels of speed, strength, endurance and flexibility, as well as shorter reaction times and proper technique have achieved excellent results.

Rarely have several success factors in elite taekwondo competitors been analysed simultaneously (in terms of medals they have or have not won at the highest levels of competition). In the above-mentioned studies, researchers focused entirely on single factors that were somatic or physiological.

The aim of this study was to identify selected success factors of elite Olympic taekwondo competitors within the context of medals that they have or have not won during Polish Junior Championships (PJCcs).

Material and methods

Sixty-three male taekwondo juniors took part in the study. The competitors were 15.90.84 years old, and their training experience was 5.11.59 years. The subjects were assigned to two groups. The first group consisted of competitors (n=27) who had won medals (MW) at PJCcs, while the other included competitors (n=36) who had finished 4th-8th (NMW) at PJCcs.

The number of points scored in the 2010-2011 PJC qualifying tournaments and at the 2010-2011 PJCcs constituted the results. The grading scale of the Polish Taekwondo Federation was employed [16].

The research included measurements of structure (A1-12), conditioning (B1-5) and flexibility (B6, 7) as well as technical (C1-3), coordination (D1-5) and energetic (E1-5) measurements.

The variables of body build assessment included body height and mass; the Quetelet index II; length of upper and lower limbs; shoulder and pelvis width; arm, forearm, thigh and shank circumference; and the thickness of five skinfolds (calf, supraspinal, tricep, subscapular, abdominal).

Conditioning was assessed using the following tests: overhead medicine ball throw (2 kg), standing broad jump, 30 s sit-ups, flying 30 m run, 20 m shuttle run (beep test) [17; 18].

Flexibility was measured with the length of front and side splits [15].

Technical skills were assessed with the time needed to perform 10 repetitions of *dollyo chagi*, *jirugii momtong*, and *360° dollyo chagi momtong* [16].

Simple and complex reactions were measured using RT (version S1) and DT (version S1) of the Vienna Test System [19].

The anaerobic muscle capacity of the lower limbs was measured using a 30 s Wingate test [20]. The following indicators were determined: workload (J·kg), peak power (W·kg), anaerobic fatigue (%) and the time of achieving and maintaining peak power (s).

At first, the correspondence of the distribution of particular variables with a normal distribution was analysed. The Shapiro-Wilk test was employed. Then, specific variables were standardised into means and standard deviations of the NMW group. The success factors were modelled using two methods. In the first—variance analysis—the scalar variable was the dichotomy of “has won/has not won a medal”. The other method incorporated modelling with the use of the general linear model (GLM)—multinomial logistic regression analysis—where the scalar variable was the number of points scored at PJC and two qualifying tournaments.

The Akaike information criterion (AIC) was adopted as the main criterion. All the available models of multinomial logistic regression (63) were analysed, and the one for which the AIC achieved minimal values was selected. Moreover, odds ratios (ORs) were calculated for particular (significant) variables incorporated in the model. The significance of the model was verified using the Wald test. The script defining an algorithm was created and was based on the P. Bieck programme [21], which was activated and performed on the R platform [22].

Results

The differences between the MW and NMW groups were noted in conditioning, in the techniques of kicks and in certain structural and coordination factors ($p < 0.05$).

The comparison of body build between the MW and NMW groups revealed that significant differences were found only in adiposity – the sum of skinfolds (A12: 10.8%) ($p < 0.05$) (Table 1).

The MW group performed better in the speed-strength (B2: 8.2%), endurance-strength (B3: 12.1%) and endurance (B5: 4.3%) tests ($p < 0.01$).

The subjects from the MW group also excelled in technical skills (fig. 1). The differences ranged from 4.8% to 7.4% (C1: 4.8%; C2: 7.4%; C3: 6.6%) ($p < 0.05$). In turn, the analysis of simple and complex reaction times revealed that competitors from the MW group demonstrated shorter complex reaction times (D5: 5.3%) than those from the NMW group ($p < 0.05$).

The Wingate test results did not reveal the dominance of either group, which may indicate that these factors did not play a key role in achieving success.

The aim of defining the regression model was to select significant success variables. The following variables were identified: standing broad jump ($p < 0.01$), 20 m shuttle run ($p < 0.001$), complex reaction time ($p < 0.05$), *360° dollyo chagi momtong* ($p < 0.05$) and the sum of skinfolds ($p < 0.001$). ORs were then calculated for those variables. ORs define the influence of unitary changes in the value of particular variables on the chances of scoring a point (Table 2).

Of all the identified variables, complex reaction time (16%) and *360° dollyo chagi momtong* (8.8%) had the largest influence on the number of points scored. Adiposity (the sum of skinfolds), lower limb muscle strength (standing broad jump) and aerobic endurance (20 m shuttle run) were less significant. The hypothesis of model incommutability as a whole was rejected at the level of $p < 0.001$. Therefore, it may be assumed that the model properly describes the investigated correlation.

Discussion

The aim of the study was to identify selected success factors of elite Olympic taekwondo competitors within the context of medals that they have or have not won during PJC.

Taekwondo competitors from the MW and NMW groups differed significantly in conditioning, technique, coordination and body build parameters. The MW group excelled in most tests.

To date, the majority of authors have focused on defining single factors, e.g., structure [6-10], biomechanics [7,11], physiology [13,23], coordination [13,24-26] and conditioning [12, 8,13,14], and the correlations of these with competition results. In our research, we tried to identify success factors while simultaneously considering what the competition result is conditioned by.

The analysis of somatic factors revealed that in WFT taekwondo competitors, low levels of adiposity were conducive to achieving success. Medal winners demonstrated lower levels of adiposity, which indicates that the mesomorphic body type was dominant. Taaffe, Pieter [2], Pieter [3], Gao [5], Marković et al. [7], Kazemi et al. [8], Fritzsche, Raschka [9], and Pieter [27] share this opinion, as they claim that the monomorphic type is the most promising in terms of achieving successful results.

Moreover, it was observed that the competitors from the MW group manifested higher levels of speed, dynamic strength as well as strength and general endurance. These results are in line with other researchers' findings, which claim that taekwondo belongs to the group of sports in which speed play a key role in achieving success [6-8,12,28-31]. In addition, the medallists performed better in the 20 m shuttle run (B5). According to McArdle et al. [32], Toskovic et al. [6], Marković [7], Cetin et al. [33], and Thompson et al. [34], this factor does not affect performance directly but rather exerts greater influence on the effectiveness and speed of energetic recovery, which may contribute to obtaining a better result and may be why our research uncovered these results.

The differences were also observed in technical skills. The medallists demonstrated considerably higher levels of

Table 1. Differences in the levels of selected groups of factors in WTF taekwondo competitors (n = 63) who have or have not won medals (mean SD)

	Group of factors (indicator)	MW group (n = 27)	NMW group (n = 36)	Differences p
structural	A1 – body height (cm)	173.3±8.4	174.1±8.9	
	A2 – body mass (kg)	64.6±12.5	67.0±13.8	
	A3 - BMI (kg/m ²)	21.3±2.58	21.9±3.01	
	A4 – lower limb length (cm)	91.0±4.98	91.7±5.41	
	A5 – upper limb length (cm)	78.0±4.96	78.9±4.96	
	A6 – shoulder width (cm)	29.7±2.06	30.1±1.82	
	A7 – pelvis width (cm)	25.8±1.55	26.3±1.61	
	A8 – arm circumference (cm)	25.4±2.79	25.9±2.66	
	A9 – forearm circumference (cm)	22.9±2.06	23.4±1.75	
	A10 – thigh circumference (cm)	51.7±5.38	52.3±5.38	
	A11 – shank circumference (cm)	35.7±3.08	35.9±2.87	
	A12 – sum of skinfolds (mm)	42.8±11.4	47.4±11.9	*
conditioning and flexibility	B1 – overhead medicine ball throw (m)	8.69±1.71	8.72±1.83	
	B2 – standing broad jump (cm)	219.9±20.1	201.8±23.0	**
	B3 – sit-ups (number)	31.5±5.05	27.7±4.22	**
	B4 – 30 m run (s)	5.07±0.39	5.26±0.48	
	B5 – 20 m shuttle run (number)	81.0±9.58	77.5±9.79	***
	B6 – side split (cm)	84.1±7.50	82.6±8.01	
	B7 – front split (cm)	60.5±13.7	50.4±8.87	
technical	C1 - dollyo chagi (s)	4.97±0.77	5.21±0.82	
	C2 – jirugi momtong (s)	2.99±0.71	3.21±0.55	
	C3 – 360° dollyo chagi momtong (s)	12.1±1.33	12.9±1.17	*
coordination	D1 – reaction time (ms)	254.1±33.0	262.1±36.2	
	D2 – time of single movement (ms)	128.1±31.6	128.7±38.4	
	D3 – proper reactions (number)	226.1±31.0	254.4±69.3	
	D4 – improper reactions (number)	22.1±9.31	24.9±8.35	
	D5 – median of reaction time (s)	0.75±0.06	0.79±0.06	*
energetic	E1 – workload (J·kg ⁻¹)	204.2±31.4	209.0±27.8	
	E2 – peak power (W·kg ⁻¹)	8.57±1.41	8.68±1.24	
	E3 – anaerobic fatigue (%)	16.9±4.48	17.9±5.08	
	E4 – time of achieving peak power (s)	8.06±1.47	7.78±1.54	
	E5 – time of maintaining peak power (s)	3.61±1.19	3.57±1.09	

*p<0.05; **p<0.01; ***p<0.001

skill than did the non-medallists, which shows how important technical skills are in terms of results. Our findings are in line with the data gathered by other researchers [8,25,30,35].

In the case of coordination, i.e. complex reaction, significant differences were noted between groups, which was similar to previous studies. Most researchers have found high correlations between time of reaction and competition results [12,13,24,25,36].

It is worth highlighting that energetic factors did not differentiate both groups, which indicates that at this level of com-

petition these factors do not determine successes or failures.

The procedure of searching all acceptable models of regression revealed that the most substantial variables in terms of success (winning PJC medals) were complex reaction time and technical skills, followed by conditioning, speed, dynamic strength and endurance.

Similar results were obtained in the process of examining elite taekwondo competitors, where conditioning, technical, structural and energetic factors were regarded as the most important [37].




Fig. 1. Normalised values of results for particular parameters in groups of taekwondo competitors

Table 2. Results of the process of seeking an optimal model

Variable	Value of estimated parameter	Standard estimation error	Significance level	OR (%)
Standing broad jump	0.01	0.002	p < 0.01	0.1
20 m shuttle run	0.009	0.004	p < 0.001	> 0.1
Complex reaction time	- 1.78	0.72	p < 0.05	16
360° dollyo chagi momtong	- 0.09	0.04	p < 0.05	8.8
The sum of skinfolds	- 0.02	0.001	p < 0.001	1.5

Note: AIC=796.7

The high levels of technical skills manifested by kick and strike performance probably resulted from an integrated manifestation of higher levels of motor and coordination abilities in the MW group.

Conclusions

- Variables concerning coordination factors (complex reaction – 16%) and technical factors (360° *dollyo chagi mon* – 8.8%) exerted the greatest influence on point scores.

2. Our findings enabled us to identify success factors in taekwondo at a national level.

- Future research should focus on seeking success factors in athletes at different levels of competition, particularly in those who achieve success at an international level, e.g., during Olympic Games.

References

- Kalina R M. Teoria sportów walki. Warszawa: Centralny Ośrodek Sportu; 2000.
- Taaffe D, Pieter W. Physical and physiological characteristics of elite taekwondo athletes. In: Commonwealth and International Conference Proceedings, Sport Science, Auckland, New Zealand; 1990; 3: 80-88.
- Pieter W. Performance characteristics of elite taekwondo athletes. Korean Journal of Sport Science 1991; 3: 94-117.
- Olds T, Kang S. Anthropometric characteristics of adult male Korean taekwondo players. The First Olympic Taekwondo Scientific Congress Proceedings. Seoul, Korea, 2000; 69-75.
- Gao B. Research on the somatotype features of Chinese elite male taekwondo athletes. J Sport Sci 2001; 21: 58-61.
- Toskovic N, Blessing D, Williford H. Physiologic profile of recreational male and female novice and experienced tae kwon do practitioners. J Sports Med Phys Fit 2004; 44: 164-172.
- Marković G, Mišigoj-Durašković M, Trninić S. Fitness profile of elite Croatian Taekwondo athletes. Collegium Antropologicum 2005; 29: 93-99.
- Kazemi M, Waalen J, Morgan C, et al. Combat sports Special Issue Research article. A profile of Olympic taekwondo competitors. J Sport Sci Med CSSI 2006; 114-121.
- Fritzsche J, Raschka C. Body composition and the somatotype of German top taekwondo practitioners. Institute of Sports Sciences Johann Wolfgang Goethe University, Frankfurt/Main. Papers Anthropology 2008; XVII: 58-71.
- Pieter W. Talent Detection in Taekwondo Practitioners. Journal of Asian Martial Arts 2010; 19(3): 9-29.
- Wąsik J. Structure of movement of a turning technique used in the event of special techniques in Taekwon-do ITF. Arch of Budo 2009; 5: 111-115.
- Melhim A. Aerobic and anaerobic power responses to the practice of taekwon do. Brit J Sport Med 2001; 35(4): 231-234.
- Abdossaleh Z, Azadeh G, Ebrahim K, et al. A survey of the physical fitness of the male taekwondo athletes of the Iranian National Team. Physical Education and Sport 2008; 6: 21-29.
- Jagiello W, Kalina R. Properties of motor development in young judokas. J Hum Kinet 2007; 17: 113-120.
- Sadowski E. Osnovy treningu koordinacyjnego sposobności w wojowniku. Białystok: Wydział Wychowania Fizycznego; 2003.
- <http://www.pztaekwondo.pl>. 4.01.2012r.
- Pilicz S. Pomiar ogólnej sprawności fizycznej. Warszawa: AWF; 1997.
- Grabowski H, Szopa J. Eurofit. Europejski Test Sprawności Fizycznej. AWF Kraków: Wydawnictwo skryptowe; 1991. (in Polish)
- Schuhfried G. Sport Psychology. Sport test battery for diagnostics and training. A-2340 Mödling, Österreich 2001; 1-43.
- Inbar O, Bar-Or O, Skinner JS. The Wingate Anaerobic Test. Human Kinetics. Champaign 1996; 95-104.
- Biecek P. Analiza danych z programem R. Modele liniowe z efektami stałymi, losowymi i mieszanymi. Warszawa: PWN; 2011.
- R Development Core Team (2011). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0.
- Matsushige K, Hartmann K, Franchini E. Taekwondo: physiological responses and match analysis. J Strength Cond Res 2009; 23(4): 1112-1117.
- Douris P. Fitness levels of middle aged martial art practitioner. Brit J Sport Med 2004; 38: 143-147.
- Saporta J, Juan Moreno's Olympic-Style Training for Taekwondo Athletes. Journal of Asian Martial Arts 2006; 15: 39-53.
- Sadowski J, Gierczuk D. Correlations between selected coordination motor abilities and technical skills of Greco-Roman wrestlers aged 14-15. Arch Budo 2009; 5: 35-39.
- Pieter W. Body build of elite junior taekwondo athletes. Acta Kinesiologiae Universitatis Tartuensis, 2008; 13: 99-106.
- Villani R, Dal Monte N, Tomasso A, et al. Study of a set of tests on rapidity as a mean of talent research in karate. Tenth annual congress of the European college of sport science. Belgrade, Serbia, July 13-16, 2005.
- Zabukovec R, Tiidus P. Physiological and anthropometric profile of elite kickboxers. J Strength Cond Res 1995; 9: 240-242.
- Pieter W, Bercades L. Strength correlates of kicking force in young taekwondo-in. In: The second international symposium for taekwondo studies. Seoul: Daekyung Books; 2009: 247-254.
- Bujak Z. Sprawność fizyczna ćwiczących taekwon-do i jej uwarunkowania. Rzeszów. Rocznik Naukowy. Ido Ruch dla Kultury 2005; V: 34-39.
- McArdle W, Katch F, Katch V. Exercise physiology: Energy nutrition, and human performance, Fifth ed. Philadelphia: Lippincott Williams & Wilkins, 2001.
- Cetin C, Karatosun H, Baydar M, et al. A regression equation to predict true maximal oxygen consumption of taekwondo athletes using a field test. Saudi Medical Journal 2005; 26(5): 848-850.
- Thompson W, Vinuela C. Physiologic profile of Tae Kwon Do black belts. Sports Med Training Rehabili 1991; 3: 49-53.
- Bujak Z. Związek korelacyjny pomiędzy budową somatyczną, koordynacją wzrokowo-ruchową i wydolnością beztlenową a wynikami sportowymi najlepszych polskich juniorów taekwon-do. In: Kierunki doskonalenia treningu i walki sportowej - diagnostyka. Warszawa: PTNKF; 2006. str. 169-174.
- Heller J, Perić T, Dlouha R et. al. Physiological profiles of male and female taekwon-do (ITF) black belts. J Sport Sci 1998; 16: 243-249.
- Sadowski J, Gierczuk D, Miller J, et al. Success factors in elite WTF taekwondo competitors. Archives of Budo, 2012; 8(3): 141-146.

Address for correspondence:

Jerzy Sadowski

Faculty of Physical Education and Sport in Białystok,
ul. Akademicka 2, Białystok, Poland
e-mail: jerzy.sadowski@awf-bp.edu.pl

Received: 16.03.2012

Accepted: 04.05.2012

